

PRODUCT DATA SHEET

Hexadecanoic acid

Catalog number: 1014

Common Name: Palmitic acid: C16:0 Fatty acid

Source: natural, plant

Solubility: chloroform, ethanol, ethyl ether

CAS number: 57-10-3

Molecular Formula: C₁₆H₃₂O₂


Molecular Weight: 256

Storage: room temperature

Purity: TLC: 99%, GC >99%

TLC System: hexane/ethyl ether/acetic acid
(85:15:1 by vol.)

Appearance: solid



Application Notes:

This high purity fatty acid is ideal as a standard and for biological studies. Hexadecanoic acid is considered the most abundant saturated fatty acid in nature comprising 20-30% of the lipids in many animal tissues. Palmitic acid has been found to cause reduced insulin activity due to its mediation of PKC- θ activation in the central nervous system.¹ During the metabolism of hexadecanoic acid it is converted to the *omega*-hydroxy hexadecanoic acid and then to the dicarboxylic hexadecanedioic acid. Long chain fatty acids have been found to inhibit the double-stranded DNA binding activity of p53 DNA binding domain suggesting that fatty acids in the cell membrane might regulate the activity of p53 for cell division, cell-cycle checkpoint, and tumor suppression.² Saturated fatty acids, such as palmitic acid, induce apoptosis in *beta*-cells which can lead to the development of diabetes.³ Long chain fatty acids acylated to sphingolipids are critical in many biological functions and substantial amounts are found to be amide-linked to the long-chain sphingoid base sphinganine, forming a ceramide, which constitutes the lipid backbone of sphingomyelin and other sphingolipids. Long chain fatty acids can often be found in esterified linkages with cholesterol, gangliosides, galactocerebrosides, sphingomyelin, and phosphatidylcholine. Hexadecanoic acid, as a saturated fatty acid, has been found to cause moderate risk of coronary heart disease as compared with polyunsaturated fatty acids but does not change the total cholesterol/high density lipoprotein-cholesterol ratio as compared with carbohydrates.⁴ Saturated fatty acids can also lead to the development of atherosclerosis in obesity and type 2 diabetes.

Selected References:

1. S. Benoit et al. "Palmitic acid mediates hypothalamic insulin resistance by altering PKC- θ subcellular localization in rodents" *Journal of Clinical Investigation*, vol. 119 pp. 2577-2589, 2009
2. H. Iijima et al. "The Inhibitory Action of Long-Chain Fatty Acids on the DNA Binding Activity of p53" *Lipids*, vol. 41 pp. 521-527, 2006
3. H. Liang et al. "Palmitic acid-induced apoptosis in pancreatic β -cells is increased by liver X receptor agonist and attenuated by eicosapentaenoate" *In Vivo*, vol. 25 pp. 711-718, 2011
4. R. Micha and D. Mozaffarian "Saturated Fat and Cardiometabolic Risk Factors, Coronary Heart Disease, Stroke, and Diabetes: a Fresh Look at the Evidence" *Lipids*, vol. 45 pp. 893-905, 2010

This product is to be used for research only. It is not intended for drug or diagnostic use, human consumption or to be used in food or food additives. Matreya assumes no liability for any use of this product by the end user. We believe the information, offered in good faith, is accurate.