

PRODUCT DATA SHEET

1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine

Catalog number: 1435

Common Name: DPPE

Source: synthetic

Solubility: chloroform/acetic acid, (95:5 by vol.);
chloroform/methanol/DI water/
acetic acid, (100:30:10:2.5 by vol.)

CAS number: 923-61-5

Molecular Formula: C₃₇H₇₄NO₈P


Molecular Weight: 692

Storage: -20°C

Purity: TLC >98%

TLC System: chloroform/methanol/DI water,
(65:25:4 by vol.)

Appearance: solid



Application Notes:

1,2-Dipalmitoyl-sn-glycero-3-phosphorylethanolamine is a glycerophospholipid in which a phosphorylethanolamine moiety occupies a terminal glycerol substitution site and palmitic acid occupies the other two substitution sites. Like most phospholipids PE usually has a saturated fatty acid on C-1 and an unsaturated fatty acid on C-2 of the glycerol backbone but the fatty acid distribution at the C-1 and C-2 positions of glycerol within all phospholipids is continually changing, owing to phospholipid degradation and the continuous phospholipid remodeling that occurs while these molecules are in membranes. PEs are neutral zwitterions at physiological pH. PE is frequently the main lipid component of microbial membranes and the second most abundant phospholipid in mammals, comprising as much as 45% of brain lipids. They are concentrated in mitochondria and are key building blocks of membrane bilayers where they are distributed asymmetrically with the majority confined to the inner leaflet. It appears that a primary role for PE, in bacterial membranes at least, is simply to dilute the high negative charge density of the anionic phospholipids. PE acts as a chaperone in transport membrane folding.¹ In animals PE is involved in the secretion of very-low-density lipoproteins and aids in membrane fusion and fission.² In plants *lyso* PE retards senescence by inhibiting phospholipase D. PE is the precursor to many important lipids, acts as a protein transport from the membrane to the vacuole, and is synthesized through the CDP-ethanolamine or the PS decarboxylation pathway. After being converted to diacyl glycerol PE acts as a second messenger.³

Selected References:

1. M. Bogdanov, W. Dowhan, "Lipid-assisted Protein Folding" *Journal of Biological Chemistry*, Vol. 274 pp. 36827-36830, 1999
2. J. Vance, "Phosphatidylserine and phosphatidylethanolamine in mammalian cells: two metabolically related aminophospholipids" *Journal of Lipid Research*, Vol. 49 pp. 1377-1387, 2008
3. D. Lang et al., "Molecular Species Analysis of 1,2-Diglycerides on Phorbol Ester Stimulation of LA-N-1 Neuroblastoma Cells During Proliferation and Differentiation" *Journal of Neurochemistry*, Vol. 65 pp. 810, 1995

This product is to be used for research only. It is not intended for drug or diagnostic use, human consumption or to be used in food or food additives. Matreya assumes no liability for any use of this product by the end user. We believe the information, offered in good faith, is accurate.